

Information-Theoretic Comparison of Quantum Many-Body Systems

K. Ch. Chatzisavvas*, C. P. Panos†, S. E. Massen‡

*Physics Department,
Aristotle University of Thessaloniki,
54124 Thessaloniki, Greece*

May, 2003

Abstract

An information measure inspired by Onicescu's information energy and Uffink's information measure (recently discussed by Brukner and Zeilinger) are calculated as functions of the number of particles N for fermionic systems (nuclei and atomic clusters) and correlated bosonic systems (atoms in a trap). Our results are compared with previous ones obtained for Shannon's information entropy, where a universal property was derived for atoms, nuclei, atomic clusters and correlated bosons. It is indicated that Onicescu's and Uffink's definitions are finer measures of information entropy than Shannon's.

Onicescu [1] introduced the concept of information energy E as a finer measure of dispersion distributions than that of Shannon's information entropy [2, 3]. So far, only the mathematical aspects of this concept have been developed, while the physical aspects have been neglected [4].

*e-mail: kchatz@auth.gr

†e-mail: chpanos@auth.gr

‡e-mail: Massen@physics.auth.gr

The information energy for a single statistical variable x with the normalized density $\rho(x)$ is defined by

$$E(\rho) = \int \rho^2(x) dx \quad (1)$$

For a Gaussian distribution of mean value μ , standard deviation σ and normalized density

$$\rho(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}} \quad (2)$$

relation (1) gives

$$E = \frac{1}{2\pi\sigma^2} \int_{-\infty}^{\infty} e^{-\frac{(x-\mu)^2}{\sigma^2}} dx$$

Thus

$$E = \frac{1}{2\sigma\sqrt{\pi}} \quad (3)$$

Therefore, the greater the information energy E , the narrower the Gaussian distribution. E does not have the dimension of energy, but it has been connected with Planck's constant appearing in Heisenberg's uncertainty relation [4, 5].

For a 3-dimensional spherically symmetric density distribution $\rho(r)$ the obvious generalization of (1) is

$$E_r = \int \rho^2(r) 4\pi r^2 dr \quad (4)$$

and

$$E_k = \int n^2(k) 4\pi k^2 dk \quad (5)$$

in position- and momentum-space respectively, where $n(k)$ is the density distribution in momentum-space.

E_r has the dimension of inverse volume, while E_k of volume. Thus the product $E_r E_k$ is dimensionless and is a measure of the concentration (or the information content) of the density distribution of a quantum system. As seen from (3) E increases as σ decreases (or the concentration increases) and Shannon's information entropy (or uncertainty) S decreases. Clearly, Shannon's information S and information energy E are reciprocal. In order to be able to compare them, we define the quantity

$$S_E = \frac{1}{E_r E_k} \quad (6)$$

as a measure of the information content of a quantum system in both position and momentum spaces.

In place of Shannon information, Brukner and Zeilinger [6] propose the quantity

$$I = \mathcal{N} \sum_{i=1}^n \left(p_i - \frac{1}{n}\right)^2 \quad (7)$$

from which they derive their notion of information content of a discrete probability distribution $p_1, p_2, \dots, p_n$. The quantity $\sum_{i=1}^n (p_i - \frac{1}{n})^2$ is one of the class of measures of the concentration of a probability distribution given by Uffink [7, 8]. For a continuous 3-dimensional density distribution $\rho(r)$, relation (7) is extended as ($\mathcal{N} = 1$)

$$I_r = \int \left(\rho(r) - \tilde{\rho}(r)\right)^2 4\pi r^2 dr \quad (8)$$

and

$$I_k = \int \left(n(k) - \tilde{n}(k)\right)^2 4\pi k^2 dk \quad (9)$$

in position- and momentum space respectively, $\tilde{\rho}(r)$ is the equivalent uniform distribution defined according to the relation

$$\tilde{\rho}(r) = \begin{cases} \rho_0 & 0 < r < R_U \\ 0 & r > R_U \end{cases} \quad (10)$$

where $\rho_0 = \text{constant}$ and $R_U = R_{\text{uniform}}$ are fixed by the relation

$$\langle r^2 \rangle_U = \langle r^2 \rangle_{\rho(r)} \quad (11)$$

where

$$\langle r^2 \rangle_U = \int_0^{R_U} \rho_0 r^2 4\pi r^2 dr \quad (12)$$

and

$$\langle r^2 \rangle_{\rho(r)} = \int_0^\infty \rho(r) r^2 4\pi r^2 dr \quad (13)$$

while

$$R_U = \sqrt{\frac{5}{3} \langle r^2 \rangle_{\rho(r)}} \quad (14)$$

and

$$\rho_0 = \frac{3}{4\pi R_U^3} \quad (15)$$

$\tilde{n}(k)$ is the equivalent uniform distribution in momentum-space, defined in a similar way. Thus we define a measure of information content by the relation

$$S_I = \frac{1}{I_r I_k} \quad (16)$$

which gives (6) putting $\tilde{\rho}(r) = \tilde{n}(k) = 0$.

We calculate S_E and S_I as functions of the number of particles N for three quantum many-body systems, where $\rho(r)$ and $\eta(k)$ are calculated numerically:

1. Nuclei, using the Skyrme III parametrization of the nuclear field [9]. Here N is the number of nucleons in nuclei.
2. Atomic clusters, employing a Woods-Saxon potential parametrized by Ekardt [10]. Here N is the number of valence electrons.
3. A correlated bosonic system (atoms in a trap) [11, 12]. Here N is the number of atoms in the trap.

In Fig.1 we plot S_E as a function of N for nuclei and clusters and in Fig.2 $S_I(N)$ for the same systems. In Fig.3 we plot $S_E(N)$ and in Fig.4 $S_I(N)$ for a correlated bosonic system. It is seen that S_E depends linearly on N for both nuclei and atomic clusters. Also S_I shows a similar trend (a power of N) for nuclei and clusters. However the dependence $S_E(N)$ and $S_I(N)$ is different for correlated bosons compared with nuclei and clusters.

Our fitted expressions are:

$$S_E(\text{clusters}) = 143.420 N, \quad S_E(\text{nuclei}) = 73.883 N \quad (\text{Fig.1})$$

$$S_I(\text{clusters}) = 431.576 N^{1.719}, \quad S_I(\text{nuclei}) = 260.275 N^{1.554} \quad (\text{Fig.2})$$

We can compare with the universal relation $S(N) = a + b \ln N$ (a, b are constants depending on the system) obtained recently [13] for Shannon's information entropy for fermionic systems (atoms, nuclei and atomic clusters) and correlated bosonic systems [11] (atoms in a trap). It was seen [13] that $S(N)$ shows the same dependence on N for all the systems considered i.e. nuclei, clusters, atoms and correlated bosons.

It is conjectured that nuclei and atomic clusters are equivalent from an information-theoretic point of view in the following sense: under any definition of information content (e.g. Shannon, Onicescu or Uffink), the dependence of information shows a similar trend (linear on $\ln N$ for Shannon, linear on N for Onicescu and a power of N for Uffink). However, the similarity breaks down for bosons. This indicates that S_E and S_I distinguish between fermions and correlated bosons i.e. they are finer measures of information than Shannon's S . Our results may contribute to the recent debate between Brukner-Zeilinger and Timpson for a possible inadequacy of the Shannon information [6, 14]

References

- [1] O. Onicescu, *C. R. Acad. Sci. Paris A* **263** (1966)
- [2] C. E. Shannon, *Bell Syst. Tech.* **27** (1948), 379; *ibid.* **27** (1948) 623
- [3] C. E. Shannon, W. Weaver, *The Mathematical Theory of Communication*, University of Illinois Press, Urbana and Chicago, 1963
- [4] M. Agop, C. Buzea, C. Gh. Buzea, L. Chirila, S. Oancea, *Chaos, Solitons and Fractals* **7** (1996), 659
- [5] H. Ioannidou, *Int. Journ. of Theor. Phys.* **20** (1981), 1129
- [6] C. Brukner, A. Zeilinger, *Phys. Rev. A* **63** (2001), 022113
- [7] J. Uffink, *PhD Thesis*, University of Utrecht (1990)
- [8] H. Maassen, J. Uffink, *Phys. Rev. Lett.* **60** (1988), 1103
- [9] C. B. Dover, N. Van Giai, *Nucl. Phys. A* **190** (1972), 373
- [10] N. Ekardt, *Phys. Rev. B* **29** (1984), 1558
- [11] S. E. Massen, Ch. C. Moustakidis, C. P. Panos, *Phys. Lett. A* **299** (2002), 131
- [12] A. Fabrocini, A. Polls, *Phys. Rev. A* **60** (1999), 2319
- [13] S. E. Massen, C. P. Panos, *Phys. Lett. A* **246** (1998), 530

[14] C. G. Timpson, LANL archive, quant-ph/0112178




Figure 1: Oniscu's information entropy S_E as function of N & for nuclei (circles) and atomic clusters (squares)




Figure 2: The same as in Fig.1 but for Uffink's information & entropy S_I




Figure 3: Dependence of S_E on N for correlated bosons




Figure 4: The same as in Fig.3 but for S_I